

LEFSCHETZ FIBRATION STRUCTURES ON KNOT SURGERY 4-MANIFOLDS

JONGIL PARK AND KI-HEON YUN

ABSTRACT. In this article we study Lefschetz fibration structures on knot surgery 4-manifolds obtained from an elliptic surface $E(2)$ using Kanenobu knots K . As a result, we get an infinite family of simply connected mutually diffeomorphic 4-manifolds coming from a pair of inequivalent Kanenobu knots. We also obtain an infinite family of simply connected symplectic 4-manifolds, each of which admits more than one inequivalent Lefschetz fibration structures of the same generic fiber.

1. INTRODUCTION

Since Seiberg-Witten theory was introduced in 1994, many techniques in 4-dimensional topology have been developed to show that a large class of simply connected smooth 4-manifolds admit infinitely many distinct smooth structures. Among them, a knot surgery technique introduced by R. Fintushel and R. Stern turned out to be one of the most powerful tools changing the smooth structure on a given 4-manifold [3]. The knot surgery construction is following: Suppose that X is a simply connected smooth 4-manifold containing an embedded torus T of square 0. Then, for any knot $K \subset S^3$, one can construct a new 4-manifold, called a *knot surgery 4-manifold*,

$$X_K = X \#_{T=T_m} (S^1 \times M_K)$$

by taking a fiber sum along a torus T in X and $T_m = S^1 \times m$ in $S^1 \times M_K$, where M_K is the 3-manifold obtained by doing 0-framed surgery along K and m is the meridian of K . Then Fintushel and Stern proved that, under a mild condition on X and T , the knot surgery 4-manifold X_K is homeomorphic, but not diffeomorphic, to a given X [3]. Furthermore, if X is a simply connected elliptic surface $E(2)$, T is the elliptic fiber, and K is a fibred knot, then it is also known that the knot surgery 4-manifold $E(2)_K$ admits not only a symplectic structure but also a genus $2g(K)+1$ Lefschetz fibration structure [5, 22]. Note that there are only two inequivalent genus one fibred knots, but there are infinitely many inequivalent genus g fibred knots for $g \geq 2$. So one may dig out some interesting properties of $E(2)_K$ by carefully investigating genus two fibred knots and related Lefschetz fibration structures.

On the one hand, Fintushel and Stern [4] conjectured that the set of all knot surgery 4-manifolds of the form $E(2)_K$ up to diffeomorphism is one-to-one correspondence with the set of all knots in S^3 up to knot equivalence. Some progresses related to the conjecture were obtained by S. Akbulut [2] and M. Akaho [1]. But a

Date: June 27, 2009.

2000 Mathematics Subject Classification. 57N13, 57R17, 53D35.

Key words and phrases. Kanenobu knot, knot surgery 4-manifold, Lefschetz fibration, monodromy factorization.

complete answer to the conjecture for prime knots up to mirror image is not known yet. Furthermore, Fintushel and Stern [5] also questioned whether any two in the following 4-manifolds

$\{Y(2; K_1, K_2) := E(2)_{K_1} \#_{id: \Sigma_{2g+1} \rightarrow \Sigma_{2g+1}} E(2)_{K_2} \mid K_1, K_2 \text{ are genus } g \text{ fibred knots}\}$ are mutually diffeomorphic or not. The second author obtained a partial result related to this question under the constraint that one of K_i ($i=1, 2$) is fixed [22].

In this article we investigate Lefschetz fibration structures on the knot surgery 4-manifold $E(2)_K$, where K ranges a family of Kanenobu knots. Remind that Kanenobu [12, 13] found an interesting family of inequivalent genus 2 fibred prime knots

$$\{K_{p,q} \mid (p, q) \in \mathcal{R}\} \text{ and } \mathcal{R} = \{(p, q) \in \mathbb{Z}^2 \mid p \in \mathbb{Z}^+, -p \leq q \leq p\},$$

where any two of them are not in mirror relation and all of them have the same Alexander polynomials. In Section 3 we consider the following family of simply connected symplectic 4-manifolds which have the same Seiberg-Witten invariants

$$\{Y(2; K_{p,q}, K_{r,s}) := E(2)_{K_{p,q}} \#_{id: \Sigma_5 \rightarrow \Sigma_5} E(2)_{K_{r,s}} \mid (p, q), (r, s) \in \mathcal{R}\}.$$

By investigating the monodromy factorization expression corresponding to Lefschetz fibration structure on $Y(2; K_{p,q}, K_{r,s})$, we answer the question raised in [5].

Theorem 1.1. *Any two simply connected symplectic 4-manifolds in*

$$\{Y(2; K_{p,q}, K_{p+1,q}) \mid p, q \in \mathbb{Z}\} \cup \{Y(2; K_{p,q}, K_{p,q+1}) \mid p, q \in \mathbb{Z}\}$$

are mutually diffeomorphic.

In section 4 we also study nonisomorphic Lefschetz fibration structures on simply connected symplectic 4-manifolds which share the same Seiberg-Witten invariants. Let $\xi_{p,q}$ be a genus five Lefschetz fibration structure on $E(2)_{K_{p,q}}$. Then, by investigating the monodromy group $G_F(\xi_{p,q})$ of $\xi_{p,q}$, we get the following theorem.

Theorem 1.2. $\xi_{p,q}$ is not equivalent to $\xi_{r,s}$ if $(p, q) \not\equiv (r, s) \pmod{2}$.

As a corollary, we can easily recapture a similar result in [18]. Remind that we constructed a pair of nonisomorphic Lefschetz fibration structures on $E(n)_K$ for a special type of 2-bridge knot K in [18]. Theorem 1.2 above also confirms such a phenomena. That is, for any $(p, q) \in \mathbb{Z}^2$ with $p \not\equiv q \pmod{2}$, $K_{p,q}$ is equivalent to $K_{q,p}$ and therefore $E(2)_{K_{p,q}}$ is the same symplectic 4-manifold as $E(2)_{K_{q,p}}$. But the theorem above implies that the corresponding Lefschetz fibration structures $\xi_{p,q}$ and $\xi_{q,p}$ are not equivalent.

Acknowledgment. Jongil Park holds a joint appointment at KIAS and in the Research Institute of Mathematics, SNU. Ki-Heon Yun was supported by Sungshin Women's University Research Grant of 2008.

2. PRELIMINARIES

In this section we briefly review some well-known facts about Lefschetz fibrations on 4-manifolds and surface mapping class groups (refer to [7] for details).

Definition 2.1. Let X be a compact, oriented smooth 4-manifold. A Lefschetz fibration is a proper smooth map $\pi : X \rightarrow B$, where B is a compact connected oriented surface and $\pi^{-1}(\partial B) = \partial X$ such that

- (1) the set of critical points $C = \{p_1, p_2, \dots, p_n\}$ of π is non-empty and lies in $\text{int}(X)$ and π is injective on C
- (2) for each p_i and $b_i := \pi(p_i)$, there are local complex coordinate charts agreeing with the orientations of X and B such that π can be expressed as $\pi(z_1, z_2) = z_1^2 + z_2^2$.

It is known that there is one-to-one correspondence between the set of symplectic Lefschetz fibrations over S^2 and the set of factorizations of the identity elements in the mapping class group as a product of right-handed Dehn twists up to Hurwitz moves and global conjugation [7, 14, 17]. Therefore a monodromy factorization of a given Lefschetz fibration has lots of information about the underlying symplectic 4-manifold.

Two monodromy factorizations W_1 and W_2 are called *Hurwitz equivalence* if W_1 can be changed to W_2 in finitely many steps of the following two operations:

- (1) *Hurwitz move*: $t_{c_n} \cdot \dots \cdot t_{c_{i+1}} \cdot t_{c_i} \cdot \dots \cdot t_{c_1} \sim t_{c_n} \cdot \dots \cdot t_{c_{i+1}}(t_{c_i}) \cdot t_{c_{i+1}} \cdot \dots \cdot t_{c_1}$
- (2) *inverse Hurwitz move*: $t_{c_n} \cdot \dots \cdot t_{c_{i+1}} \cdot t_{c_i} \cdot \dots \cdot t_{c_1} \sim t_{c_n} \cdot \dots \cdot t_{c_i} \cdot t_{c_i}^{-1}(t_{c_{i+1}}) \cdot \dots \cdot t_{c_1}$

where $t_a(t_b) = t_{t_a(b)}$ and it is $t_a \circ t_b \circ t_a^{-1}$ as an element of mapping class group. This relation comes from the choice of Hurwitz system, a set of mutually disjoint arcs except the base point b_0 which connecting b_0 to b_i .

A choice of generic fiber also gives another equivalence relation. Two monodromy factorizations W_1 and W_2 are called *simultaneous conjugation equivalence* if $W_2 = f(W_1)$ for some $f \in \mathcal{M}_g$, where Σ_g is a generic fiber of the Lefschetz fibration W_1 .

Two Lefschetz fibrations $f_1 : X_1 \rightarrow B_1$, $f_2 : X_2 \rightarrow B_2$ are called *isomorphic* if there are orientation preserving diffeomorphisms $H : X_1 \rightarrow X_2$ and $h : B_1 \rightarrow B_2$ such that the following diagram commutes:


$$(2.1) \quad \begin{array}{ccc} X_1 & \xrightarrow{H} & X_2 \\ f_1 \downarrow & & \downarrow f_2 \\ B_1 & \xrightarrow{h} & B_2 \end{array}$$

Monodromy factorizations of two isomorphic Lefschetz fibrations are related by a sequence of Hurwitz equivalences and simultaneous conjugation equivalences.

Notation. We denote by $W_1 \cong W_2$ if two monodromy factorizations W_1 and W_2 are equivalent. In the case that two manifolds X_1 and X_2 are diffeomorphic, we denote by $X_1 \approx X_2$.

Definition 2.2. Let $\pi : X \rightarrow S^2$ be a Lefschetz fibration and let F be a fixed generic fiber of the Lefschetz fibration. Let $W = w_n \cdot \dots \cdot w_2 \cdot w_1$ be a monodromy factorization of the Lefschetz fibration corresponding to F . Then the *monodromy group* $G_F(W)$ is a subgroup of the mapping class group $\mathcal{M}_F = \pi_0(\text{Diff}^+(F))$ generated by $w_1, w_2, \dots, w_n$. We will write $G(W)$ when the generic fiber F is clear from context. The element $w_n \circ \dots \circ w_2 \circ w_1$ in $\mathcal{M}_F$ is denoted by λ_W .

Lemma 2.3. *If two monodromy factorizations W_1 and W_2 give isomorphic Lefschetz fibrations over S^2 , then monodromy groups $G(W_1)$ and $G(W_2)$ are isomorphic as a subgroup of the mapping class group $\mathcal{M}_F$. Moreover if we fix a generic fiber F , then $G_F(W_1) = G_F(W_2)$.*

FIGURE 1. an involution and its vanishing cycles with $g = 2$

A monodromy factorization of a Lefschetz fibration structure on $E(n)_K$ was studied by Fintushel and Stern [5] and we could find an explicit monodromy factorization of $E(n)_K$ [22] with the help of factorization of the identity element in the mapping class group which were discovered by Y. Matsumoto [17], M. Korkmaz [16] and Y. Gurtas [8].

Definition 2.4. Let $M(n, g)$ be the desingularization of the double cover of $\Sigma_g \times S^2$ branched over $2n(\{pt.\} \times S^2) \cup 2(\Sigma_g \times \{pt.\})$.

Lemma 2.5 ([16, 21]). $M(2, g)$ has a monodromy factorization $\eta_{1,g}^2$, where

$$\eta_{1,g} = t_{B_0} \cdot t_{B_1} \cdot t_{B_2} \cdots t_{B_{2g}} \cdot t_{B_{2g+1}} \cdot t_{b_{g+1}}^2 \cdot t_{b'_{g+1}}^2$$

and $\{B_j, b_{g+1}, b'_{g+1}\}$ are simple closed curves on Σ_{2g+1} as in Figure 1.

Theorem 2.6 ([5, 22]). Let $K \subset S^3$ be a fibred knot of genus g . Then $E(2)_K$, as a genus $(2g + 1)$ Lefschetz fibration, has a monodromy factorization of the form

$$\Phi_K(\eta_{1,g}) \cdot \Phi_K(\eta_{1,g}) \cdot \eta_{1,g} \cdot \eta_{1,g},$$

where $\eta_{1,g}^2$ is a monodromy factorization of $M(2, g)$ and

$$\Phi_K = \varphi_K \oplus id \oplus id : \Sigma_g \# \Sigma_1 \# \Sigma_g \rightarrow \Sigma_g \# \Sigma_1 \# \Sigma_g$$

is a diffeomorphism obtained by using a (geometric) monodromy φ_K of K defined by

$$S^3 \setminus \nu(K) = (I \times \Sigma_g^1) / ((1, x) \sim (0, \varphi_K(x))),$$

where Σ_g^1 is an oriented surface of genus g with one boundary component.

3. ISOMORPHIC LEFSCHETZ FIBRATIONS

In this section we construct examples of simply connected isomorphic symplectic Lefschetz fibrations with the same generic fiber but coming from a pair of inequivalent fibred knots. In [5] Fintushel and Stern constructed families of simply connected symplectic 4-manifolds with the same Seiberg-Witten invariants. Among them, they considered a set of the following symplectic 4-manifolds

$$\{Y(2; K_1, K_2) := E(2)_{K_1} \#_{id: \Sigma_{2g+1} \rightarrow \Sigma_{2g+1}} E(2)_{K_2} \mid K_1, K_2 \text{ are genus } g \text{ fibred knots}\}$$

and they showed that

$$SW_{Y(2; K_1, K_2)} = t_K + t_K^{-1}.$$

In [22] we found examples such that $Y(2; K, K_1)$ and $Y(2; K, K_2)$ are diffeomorphic even though K_1 is not equivalent to K_2 . In this section we will generalize such a

construction. That is, we will construct infinitely many pairs (K, K') of inequivalent genus 2 fibred knots such that all of $Y(2; K, K')$'s are mutually diffeomorphic.

A family of inequivalent knots with the same Alexander polynomials were constructed by several authors. Among them, Kinoshita and Terasaka [15] constructed a nontrivial knot with the trivial Alexander polynomial by using an operation, so called knot union. After that, Kanenobu constructed infinitely many inequivalent knots $K_{p,q}$ ($p, q \in \mathbb{Z}$) with the same Alexander polynomials [12, 13]. They constructed the examples from the ribbon fibred knot $4_1 \# (-4_1^*)$ by applying the Stallings' twist [20] at two different locations repeatedly, where K^* is the mirror image of K .




FIGURE 2. A Kanenobu knot $K_{p,q}$

The followings are known to Kanenobu.

Lemma 3.1 ([12]). *Let $K_{p,q}$ be a Kanenobu knot as in Figure 2. Then*

- (1) $K_{0,0} = 4_1 \# (-4_1^*)$
- (2) *The Alexander matrix of $K_{p,q}$ is*
$$\begin{pmatrix} t^2 - 3t + 1 & (p-q)t \\ 0 & t^2 - 3t + 1 \end{pmatrix}$$
- (3) $\Delta_{K_{p,q}}(t) \doteq (t - 3 + t^{-1})^2$
- (4) $K_{p,q}$ *is a fibred ribbon knot*
- (5) $K_{p,q} \sim K_{r,s}$ *if and only if* $(p, q) = (r, s) \text{ or } (s, r)$
- (6) $K_{p,q}^* \sim K_{-q, -p}$
- (7) $K_{p,q}$ *is a prime knot if* $(p, q) \neq (0, 0)$




FIGURE 3. Standard simple closed curves

It is not hard to see [9] that the monodromy map $\Phi_{K_{p,q}}$ of a Kanenobu knot $K_{p,q}$ is

$$t_d^q \circ t_{c_2}^p \circ t_{a_2} \circ t_{b_2}^{-1} \circ t_{a_1}^{-1} \circ t_{b_1},$$

where $\{a_i, b_i, c_i, d\}$ are simple closed curves in Figure 3. Therefore we get that $Y(2; K_{p,q}, K_{r,s})$ has a monodromy factorization of the form

$$\Phi_{K_{r,s}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2 \cdot \Phi_{K_{p,q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2.$$

Lemma 3.2. *For each $k \in \mathbb{Z}_{\geq 0}$ and fixed p, q , we have following relations:*

$$(3.1) \quad t_{c_2} = \lambda_{t_{c_2}^k(t_{B_2})}(t_{c_2}^{k+1}(t_{B_2})) = \lambda_{t_{c_2}^{k+1}(t_{B_3})}(t_{c_2}^k(t_{B_3}))$$

$$(3.2) \quad t_d = \lambda_{t_d^k(t_{B_4})}(t_d^{k+1}(t_{B_4})) = \lambda_{t_d^{k+1}(t_{B_3})}(t_d^k(t_{B_3}))$$

$$(3.3) \quad t_{c_2} = \lambda_{\Phi_{K_{k,q}}(t_{B_3})}(\Phi_{K_{k+1,q}}(t_{B_3}))$$

$$(3.4) \quad t_d = \lambda_{\Phi_{K_{p,k+1}}(t_{B_4})}^{-1}(\Phi_{K_{p,k}}(t_{B_4})).$$

Therefore

$$\begin{aligned} t_{c_2} &\in G_F(\xi_{p,q} \cdot \xi_{p\pm 1,q}), & t_{c_2} &\in G_F(\eta_{1,2}^2 \cdot t_{c_2}(\eta_{1,2}^2)) \\ t_d &\in G_F(\xi_{p,q} \cdot \xi_{p,q\pm 1}), & t_d &\in G_F(\eta_{1,2}^2 \cdot t_d(\eta_{1,2}^2)). \end{aligned}$$

Proof. It is easy to see the image of a simple closed curve on an oriented surface Σ_5 under Dehn twists and we get

$$(3.5) \quad c_2 = (t_{B_2} \circ t_{c_2})(B_2) = (t_{B_3}^{-1} \circ t_{c_2}^{-1})(B_3)$$

$$(3.6) \quad d = (t_{B_4} \circ t_d)(B_4) = (t_{B_3} \circ t_d)(B_3).$$

Equation (3.1) is easily obtained from Equation (3.5) because

$$(t_{B_2} \circ t_{c_2})(t_{B_2}) = t_{c_2} = (t_{B_3}^{-1} \circ t_{c_2}^{-1})(t_{B_3})$$

and, from this, we obtain

$$\begin{aligned} t_{c_2} = t_{t_{c_2}^k(c_2)} &= \lambda_{t_{c_2}^k(t_{c_2})} \\ &= \lambda_{t_{c_2}^k(\lambda_{t_{B_2} \circ t_{c_2}}(t_{B_2}))} \\ &= \lambda_{t_{c_2}^k \circ t_{B_2} \circ t_{c_2}}(t_{B_2}) \\ &= \lambda_{t_{c_2}^k \circ t_{B_2} \circ t_{c_2}^{-k} \circ t_{c_2}^{k+1}}(t_{B_2}) \\ &= \lambda_{t_{c_2}^k(t_{B_2})}(t_{c_2}^{k+1}(t_{B_2})) \end{aligned}$$

and

$$\begin{aligned} \lambda_{t_{c_2}^{k+1}(t_{B_3})}(t_{c_2}^k(t_{B_3})) &= t_{(t_{c_2}^{k+1}(t_{B_3}) \circ t_{c_2}^k)(B_3)} \\ &= t_{(t_{c_2}^{k+1} \circ t_{B_3}^{-1} \circ t_{c_2}^{-1})(B_3)} \\ &= t_{t_{c_2}^{k+1}(c_2)} = t_{c_2}. \end{aligned}$$

Similarly, we get Equation (3.2) from Equation (3.6).

Next we prove Equation (3.3). First, observe that

$$(\Phi_{K_{0,0}}(t_{B_3}) \circ t_{c_2} \circ \Phi_{K_{0,0}})(B_3) = c_2$$

because $\Phi_{K_{0,0}}(B_3)$ meets with c_2 at one point. Therefore

$$\begin{aligned} \lambda_{\Phi_{K_{k,q}}(t_{B_3})}(\Phi_{K_{k+1,q}}(t_{B_3})) &= t_{(\Phi_{K_{k,q}}(t_{B_3}) \circ \Phi_{K_{k+1,q}})(B_3)} \\ &= t_{(t_d^q \circ t_{c_2}^k \circ \Phi_{K_{0,0}}(t_{B_3}) \circ t_{c_2} \circ \Phi_{K_{0,0}})(B_3)} \\ &= t_{(t_d^q \circ t_{c_2}^p)(c_2)} = t_{c_2}. \end{aligned}$$

For equation (3.4), observe that

$$(\Phi_{K_{0,0}}(t_{B_4}^{-1}) \circ t_d^{-1} \circ \Phi_{K_{0,0}})(B_4) = d$$

and it implies

$$\begin{aligned} \lambda_{\Phi_{K_{p,k+1}}(t_{B_4})}^{-1}(\Phi_{K_{p,k}}(t_{B_4})) &= t_{(\Phi_{K_{p,k+1}}(t_{B_4}^{-1}) \circ \Phi_{K_{p,k}})(B_4)} \\ &= t_{(t_d^{k+1} \circ t_{c_2}^p \circ \Phi_{K_{0,0}}(t_{B_4}^{-1}) \circ t_d^{-1} \circ \Phi_{K_{0,0}})(B_4)} \\ &= t_{(t_d^{k+1} \circ t_{c_2}^p)(d)} = t_d. \end{aligned}$$

□

Lemma 3.3. *For each pair $p, q \in \mathbb{Z}$, we get diffeomorphisms*

$$Y(2; K_{p,q}, K_{p+1,q}) \approx Y(2; K_{p+1,q}, K_{p+2,q})$$

and

$$Y(2; K_{p,q}, K_{p,q+1}) \approx Y(2; K_{p,q+1}, K_{p,q+2}).$$

Proof. $Y(2; K_{p,q}, K_{p+1,q})$ has a monodromy factorization of the form

$$\Phi_{K_{p+1,q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2 \cdot \Phi_{K_{p,q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2$$

where $\Phi_{p,q} = t_d^q \circ t_{c_2}^p \circ t_{a_2} \circ t_{b_2}^{-1} \circ t_{a_1}^{-1} \circ t_{b_1}$.

By Equation (3.3) in Lemma 3.2,

$$\begin{aligned} t_{c_2} &\in G_F(\Phi_{K_{p+1,q}}(\eta_{1,2}^2) \cdot \Phi_{K_{p,q}}(\eta_{1,2}^2)) \\ t_{c_2} &\in G_F(t_{c_2}(\eta_{1,2}^2) \cdot \eta_{1,2}^2). \end{aligned}$$

Therefore we get

$$\begin{aligned} &\Phi_{K_{p+1,q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2 \cdot \Phi_{K_{p,q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2 \\ &\sim \Phi_{K_{p+1,q}}(\eta_{1,2}^2) \cdot \Phi_{K_{p,q}}(\eta_{1,2}^2) \cdot t_{c_2}(\eta_{1,2}^2) \cdot \eta_{1,2}^2 \\ &\sim \Phi_{K_{0,q}}(\eta_{1,2}^2) \cdot \Phi_{K_{0,q}}(\eta_{1,2}^2) \cdot t_{c_2}(\eta_{1,2}^2) \cdot \eta_{1,2}^2. \end{aligned}$$

It implies that, for each fixed q , $Y(2; K_{p,q}, K_{p+1,q})$ has isomorphic Lefschetz fibration structures, so that they are all diffeomorphic.

Similarly, by using Equation (3.4) in Lemma 3.2, we can prove that

$$Y(2; K_{p,q}, K_{p,q+1}) \approx Y(2; K_{p,q+1}, K_{p,q+2}).$$

□

Theorem 3.4. *Any two simply connected symplectic 4-manifolds in*

$$\{Y(2; K_{p,q}, K_{p+1,q}) \mid p, q \in \mathbb{Z}\} \cup \{Y(2; K_{p,q}, K_{p,q+1}) \mid p, q \in \mathbb{Z}\}$$

are mutually diffeomorphic.

Proof. By Lemma 3.3, the smooth classification problems of simply connected symplectic 4-manifolds in $\{Y(2; K_{p,q}, K_{r,s}) \mid (r = p \pm 1, s = q) \text{ or } (r = p, s = q \pm 1)\}$ can be reduced to the smooth classification problems of 4-manifolds in $\{Y(2; K_{p,p}, K_{p+1,p}), Y(2; K_{p,p}, K_{p,p+1}) \mid p \in \mathbb{Z}\}$. Then, by Lemma 3.1, we have $K_{p,p \pm 1} \sim K_{p \pm 1, p}$ and it implies

$$\begin{aligned} Y(2; K_{p,p}, K_{p+1,p}) &\approx Y(2; K_{p,p}, K_{p,p+1}) \\ Y(2; K_{p,p+1}, K_{p+1,p+1}) &\approx Y(2; K_{p+1,p}, K_{p+1,p+1}). \end{aligned}$$

Furthermore, by Lemma 3.3, we also get

$$Y(2; K_{p,p+1}, K_{p+1,p+1}) \approx Y(2; K_{p+1,p+1}, K_{p+2,p+1}).$$

Therefore, for any $p, q \in \mathbb{Z}$, we have

$$Y(2; K_{p,p}, K_{p,p\pm 1}) \approx Y(2; K_{q,q}, K_{q\pm 1,q})$$

and it implies the conclusion. $\square$

4. NONISOMORPHIC LEFSCHETZ FIBRATIONS

In this section we investigate some algebraic and graph theoretic properties of $\xi_{p,q} = \Phi_{K_{p,q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2$ and its monodromy group $G_{\Sigma_5}(\xi_{p,q})$. In [11], Humphries showed that the minimal number of Dehn twist generators of the mapping class group $\mathcal{M}_g$ or $\mathcal{M}_g^1$ is $2g+1$ by using symplectic transvection and modulo two Euler number of a graph.

Definition 4.1. Let $\{\gamma_1, \gamma_2, \dots, \gamma_{2g}\}$ be a set of simple closed curves on Σ_g which generate $H_1(\Sigma_g; \mathbb{Z}_2)$. Let $\Gamma(\gamma_1, \gamma_2, \dots, \gamma_{2g})$ be a graph which is defined by

- a vertex for each simple closed curve γ_i , $i = 1, 2, \dots, 2g$
- an edge between γ_i and γ_j if $i(\gamma_i, \gamma_j) = 1 \pmod{2}$ where $i(\gamma_i, \gamma_j)$ is the minimum number of intersection between two simple closed curves γ_i and γ_j
- no intersections between any two edges.

Let γ be a simple closed curve on Σ_5 , then $\gamma = \sum_{i=1}^{2g} \varepsilon_i \gamma_i$ ($\varepsilon_i = 0$ or 1) as an element of $H_1(\Sigma_g; \mathbb{Z}_2)$. Let $\bar{\gamma} := \cup_{\varepsilon_i=1} \bar{\gamma}_i$ where $\bar{\gamma}_i$ be the union of all closure of half edges with one end vertex γ_i . We define $\chi_\Gamma(\gamma)$ as the modulo 2 Euler number $\chi_\Gamma(\bar{\gamma})$.

Lemma 4.2. Let $\Gamma(\gamma_1, \dots, \gamma_{2g})$ be a graph of simple closed curves $\{\gamma_1, \dots, \gamma_{2g}\}$ which generate $\mathbb{Z}_2$ -vector space $H_1(\Sigma_g; \mathbb{Z}_2)$. Let $G_{\Gamma,g}$ be a subgroup of $\mathcal{M}_g$ generated by

$$\{t_\alpha \mid \alpha \text{ is a nonseparating simple closed curve on } \Sigma_g \text{ such that } \chi_\Gamma(\alpha) = 1\}.$$

Then $G_{\Gamma,g}$ is a nontrivial proper subgroup of $\mathcal{M}_g$. Moreover, if β is a non-separating simple closed curve on Σ_g with $\chi_\Gamma(\beta) = 0$, then $t_\beta \notin G_{\Gamma,g}$.

Proof. Let us prove that $G_{\Gamma,g}$ is a nontrivial proper subgroup of $\mathcal{M}_g$. Each element in $H_1(\Sigma_g; \mathbb{Z}_2)$ can be represented by a non-separating simple closed curve on Σ_g and $\mathcal{M}_g$ acts transitively on the set of all non-separating simple closed curves on Σ_g .

If c is a non-separating simple closed curve on Σ_g such that $\chi_\Gamma(c) = 1$, then

$$t_c(\gamma) = \begin{cases} \gamma, & \text{if } i(c, \gamma) = 0 \pmod{2} \\ c + \gamma, & \text{if } i(c, \gamma) = 1 \pmod{2} \end{cases}$$

and, for the $i(c, \gamma) = 1 \pmod{2}$ case,

$$\overline{t_c(\gamma)} = \bar{c} \cup \bar{\gamma} \text{ and } \bar{c} \cap \bar{\gamma} = \text{odd number of points}.$$

So

$$\chi_\Gamma(t_c(\gamma)) = \begin{cases} \chi_\Gamma(\gamma), & \text{if } i(c, \gamma) = 0 \pmod{2} \\ \chi_\Gamma(\bar{\gamma}) + \chi_\Gamma(\bar{c}) + \text{odd number} \equiv \chi_\Gamma(\gamma), & \text{if } i(c, \gamma) = 1 \pmod{2}. \end{cases}$$

For any $f \in G_{\Gamma,g}$, f is of the form $t_{c_k}^{\epsilon_k} \circ t_{c_{k-1}}^{\epsilon_{k-1}} \circ \dots \circ t_{c_2}^{\epsilon_2} \circ t_{c_1}^{\epsilon_1}$, where each c_i is a non-separating simple closed curve with $\chi_{\Gamma}(c_i) = 1$ and it implies $\chi_{\Gamma}(f(\gamma)) \equiv \chi_{\Gamma}(\gamma) \pmod{2}$. Therefore if $G_{\Gamma,g} = \mathcal{M}_g$, then for any non-separating simple closed curves γ on Σ_g we have to have $\chi_{\Gamma}(\gamma) = 1$. It is clearly impossible. So $G_{\Gamma,g}$ is a nontrivial proper subgroup of $\mathcal{M}_g$.

If β is a non-separating simple closed curve with $\chi_{\Gamma}(\beta) = 0$, then, for simple closed curve γ on Σ_g with $i(\beta, \gamma) = 1$, we have $\chi_{\Gamma}(t_{\beta}(\gamma)) \not\equiv \chi_{\Gamma}(\gamma) \pmod{2}$. Therefore $t_{\beta} \notin G_{\Gamma,g}$. $\square$

Lemma 4.3. *For any $p, q \in \mathbb{Z}$,*

$$t_{c_2}, t_d \notin G_F(\xi_{p,q}) = G_F(\Phi_{K_{p,q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2).$$

Proof. We will prove this in four cases.

Case 1: p and q are even integers: Let us consider

$$\Gamma_1 = \Gamma(\{c_1, a_1, a_2, b_2, a_3, b_3, a_4, a_5, B_2, B_4\})$$

where $\{a_i, b_i, c_i, d_i, B_i\}$ are simple closed curves on Σ_5 as in Figure 1 and in Figure 3. Then graph Γ_1 is given as in Figure 4 and we have following relations in $H_1(\Sigma_5; \mathbb{Z}_2)$:

$$(4.1) \quad \begin{aligned} b_2 &= c_1 + c_2, \\ b_3 &= c_1 + c_2 + c_3 = b'_3, \\ B_0 &= a_1 + a_2 + a_3 + a_4 + a_5, \\ B_1 &= B_2 + a_1 + a_5, \\ B_2 &= B_4 + a_2 + a_4 + c_2 + c_5, \\ B_3 &= B_4 + a_2 + a_4, \\ B_4 &= a_3 + c_3 + c_4, \\ B_5 &= a_3 + b_3 + b'_3 = a_3. \end{aligned}$$




FIGURE 4. Graph Γ_1

Therefore we get

$$\begin{aligned} c_2 &= b_2 + c_1, \\ c_3 &= b_2 + b_3, \\ c_4 &= B_4 + a_3 + b_2 + b_3, \\ c_5 &= B_2 + B_4 + a_2 + a_4 + b_2 + c_1, \\ c_6 &= c_1 + c_2 + c_3 + c_4 + c_5 = c_1 + a_2 + a_3 + a_4 + B_2 \end{aligned}$$

and it implies that each c_i is in the space spanned by

$$\{c_1, a_1, a_2, b_2, a_3, b_3, a_4, a_5, B_2, B_4\}.$$

Since $b_i = c_1 + \cdots + c_i$, each b_i is also in the space spanned by

$$\{c_1, a_1, a_2, b_2, a_3, b_3, a_4, a_5, B_2, B_4\}$$

and therefore $\{c_1, a_1, a_2, b_2, a_3, b_3, a_4, a_5, B_2, B_4\}$ generates $H_1(\Sigma_5; \mathbb{Z}_2)$.

Furthermore, by drawing figures, we can easily obtain the following relations:

$$\begin{aligned} \Phi_{K_{0,0}}(B_5) &= B_5, \\ \Phi_{K_{0,0}}(B_4) &= B_4 + a_2, \\ \Phi_{K_{0,0}}(B_3) &= B_3 + b_2 = B_4 + a_2 + a_4 + b_2, \\ \Phi_{K_{0,0}}(B_2) &= B_2 + a_1 + b_2 + a_2, \\ \Phi_{K_{0,0}}(B_1) &= B_1 + b_1 + b_2 + a_2 = B_2 + a_1 + a_2 + a_5 + b_1 + b_2, \\ \Phi_{K_{0,0}}(B_0) &= B_0 + a_1 + b_1 + a_2 + b_2 = a_3 + a_4 + a_5 + b_1 + b_2. \end{aligned}$$

Therefore

$$\chi_{\Gamma_1}(a_i) = \chi_{\Gamma_1}(B_i) = \chi_{\Gamma_1}(\Phi_{K_{0,0}}(B_i)) = 1, \text{ for } i = 0, 1, 2, 3, 4, 5$$

and $\chi_{\Gamma_1}(c_1) = \chi_{\Gamma_1}(c_6) = 1$. So we have

$$\{t_{B_i}, \Phi_{K_{0,0}}(t_{B_i}), t_{a_j}, t_{b_3}, t_{b'_3}, t_{c_1}, t_{c_6} \mid i = 0, 1, 2, 3, 4, 5, j = 1, 2, 3, 4, 5\} \subset G_{\Gamma_1, 5}$$

and each generator of the group $G_F(\Phi_{K_{0,0}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2)$ is an element of $G_{\Gamma_1, 5}$. It implies that $G_F(\Phi_{K_{0,0}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2) \leq G_{\Gamma_1, 5}$.

But we have

$$\chi_{\Gamma_1}(c_j) = \chi_{\Gamma_1}(d) = 0$$

for $j = 2, 3, 4, 5$ and therefore

$$t_{c_2}, t_{c_3}, t_{c_4}, t_{c_5}, t_d \notin G_{\Gamma_1, 5}.$$

It implies that $t_{c_2}, t_d \notin G_F(\Phi_{K_{0,0}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2)$.

Since $\mathbb{Z}_2$ -homology class of $\Phi_{K_{2p,2q}}(B_i)$ and $\Phi_{K_{0,0}}(B_i)$ are the same for any $p, q \in \mathbb{Z}$, we get

$$\chi_{\Gamma_1}(\Phi_{K_{2p,2q}}(B_i)) = \chi_{\Gamma_1}(\Phi_{K_{0,0}}(B_i))$$

for $i = 0, 1, 2, 3, 4, 5$. It implies that $G_F(\Phi_{K_{2p,2q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2) \leq G_{\Gamma_1, 5}$ and therefore we have $t_{c_2}, t_d \notin G_F(\Phi_{K_{2p,2q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2)$.



Case 2: p is an odd and q is an even integer: Let us consider

$$\Gamma_2 = \Gamma(\{a_3, b_3, B_1, B_2, B_3, B_4, d_1, d_2, d_3, d_4\})$$

where $\{a_i, b_i, c_i, d_i, B_i\}$ are simple closed curves on Σ_5 as in Figure 1, Figure 3 and Figure 5. Then the graph Γ_2 is as in Figure 6.

Since the equations in (4.1) are still valid and we have

$$\begin{aligned} d_1 &= a_1 + a_3, \\ d_2 &= a_3 + a_4, \\ d_3 &= c_2 + c_3, \\ d_4 &= c_2 + c_3 + c_5 + c_6, \\ b_3 &= c_1 + c_2 + c_3 = c_4 + c_5 + c_6, \end{aligned}$$

FIGURE 5. Simple closed curves d_i FIGURE 6. Graph Γ_2

we get

$$\begin{aligned}
 a_1 &= d_1 + a_3, \\
 a_2 &= B_3 + B_4 + a_3 + d_2, \\
 a_3 &= B_5, \\
 a_4 &= a_3 + d_2, \\
 (4.2) \quad a_5 &= B_1 + B_2 + a_3 + d_1, \\
 c_1 &= b_3 + d_3, \\
 c_2 &= a_3 + b_3 + d_4 + B_4, \\
 c_3 &= a_3 + b_3 + d_3 + d_4 + B_4, \\
 c_4 &= b_3 + d_3 + d_4, \\
 c_5 &= a_3 + b_3 + c_2 + d_4 + B_2 + B_3 + B_4.
 \end{aligned}$$

Hence Equations (4.2) implies that $\{a_3, b_3, B_1, B_2, B_3, B_4, d_1, d_2, d_3, d_4\}$ is a basis of $\mathbb{Z}_2$ -vector space $H_1(\Sigma_5; \mathbb{Z}_2)$.

Since $\Phi_{K_{1,0}} = t_{c_2} \circ t_{a_2} \circ t_{b_2}^{-1} \circ t_{a_1}^{-1} \circ t_{b_1}$, using (4.2) and $b_i = c_1 + \dots + c_i$, we get

$$\begin{aligned}
\Phi_{K_{1,0}}(B_0) &= B_0 + a_1 + b_1 + a_2 + b_2, \\
\Phi_{K_{1,0}}(B_1) &= B_1 + b_1 + b_2 + a_2 + c_2 \\
&= B_1 + B_3 + B_4 + a_3 + d_2, \\
(4.3) \quad \Phi_{K_{1,0}}(B_2) &= B_2 + a_1 + b_2 + a_2 + c_2 \\
&= B_2 + B_3 + B_4 + b_3 + d_1 + d_2 + d_3, \\
\Phi_{K_{1,0}}(B_3) &= B_3 + b_2 + c_2 = B_3 + b_3 + d_3, \\
\Phi_{K_{1,0}}(B_4) &= B_4 + a_2 + c_2 = B_3 + B_4 + b_3 + d_2 + d_4, \\
\Phi_{K_{1,0}}(B_5) &= B_5.
\end{aligned}$$

A computation of χ_{Γ_2} shows that

$$(4.4) \quad \chi_{\Gamma_2}(B_i) = \chi_{\Gamma_2}(\Phi_{K_{1,0}}(B_i)) = \chi_{\Gamma_2}(b_3) = \chi_{\Gamma_2}(b'_3) = \chi_{\Gamma_2}(a_3) = 1$$

for each $i = 0, 1, 2, 3, 4, 5$ and

$$(4.5) \quad \chi_{\Gamma_2}(c_1) = \chi_{\Gamma_2}(c_2) = \chi_{\Gamma_2}(a_1) = \chi_{\Gamma_2}(a_2) = \chi_{\Gamma_2}(b_2) = \chi_{\Gamma_2}(d) = 0.$$

Therefore $G_F(\Phi_{K_{1,0}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2) \leq G_{\Gamma_2,5}$ and, since $t_{c_2}, t_d \notin G_{\Gamma_2,5}$, we get

$$t_{c_2}, t_d \notin G_F(\Phi_{K_{1,0}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2).$$

Furthermore, since $\Phi_{K_{2p+1,2q}}(B_i)$ and $\Phi_{K_{1,0}}(B_i)$ represent the same element in $H_1(\Sigma_2; \mathbb{Z}_2)$, we get $\chi_{\Gamma_2}(\Phi_{K_{2p+1,2q}}(B_i)) = \chi_{\Gamma_2}(\Phi_{K_{1,0}}(B_i)) = 1$ and it implies that

$$t_{c_2}, t_d \notin G_F(\Phi_{K_{2p+1,2q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2)$$

for any $p, q \in \mathbb{Z}$ because $G_F(\Phi_{K_{2p+1,2q}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2) \leq G_{\Gamma_2,5}$.

Case 3: p is an even and q is an odd integer: We want to find a graph

$$\Gamma_3 = \Gamma(\{\gamma_1, \gamma_2, \dots, \gamma_{10}\})$$

satisfying

$$(4.6) \quad \chi_{\Gamma_3}(B_i) = \chi_{\Gamma_3}(\Phi_{K_{0,1}}(B_i)) = \chi_{\Gamma_3}(b_3) = \chi_{\Gamma_3}(b'_3) = \chi_{\Gamma_3}(a_3) = 1$$

for $i = 0, 1, 2, 3, 4, 5$ and

$$(4.7) \quad \chi_{\Gamma_3}(c_2) = \chi_{\Gamma_3}(d) = 0.$$

Note that we observe the following relations in $H_1(\Sigma_5; \mathbb{Z}_2)$:

$$(4.8) \quad
\begin{array}{|c|c|c|}
\hline
& \Phi_{K_{0,0}}(B_i) & \Phi_{K_{0,1}}(B_i) \\
\hline
B_0 & B_0 + a_1 + b_1 + a_2 + b_2 & B_0 + a_1 + b_1 + a_2 + b_2 \\
\hline
B_1 & B_1 + b_1 + b_2 + a_2 & B_1 + b_1 + a_2 + b_2 + d \\
\hline
B_2 & B_2 + a_1 + b_2 + a_2 & B_2 + a_1 + b_2 + a_2 \\
\hline
B_3 & B_3 + b_2 & B_3 + b_2 \\
\hline
B_4 & B_4 + a_2 & B_4 + a_2 + d \\
\hline
B_5 & B_5 & B_5 \\
\hline
\end{array}$$

Hence, by Lemma 4.2 and Equations (4.6), (4.7) and (4.8), we need to find a graph Γ_3 such that

- even number of $\{a_1, b_1, a_2, b_2\}$ have $\chi_{\Gamma_3} = 0$
- odd number of $\{b_1, b_2, a_2\}$ have $\chi_{\Gamma_3} = 0$
- even number of $\{a_1, b_2, a_2\}$ have $\chi_{\Gamma_3} = 0$

- even number of $\{b_2\}$ have $\chi_{\Gamma_3} = 0$
- odd number of $\{a_2\}$ have $\chi_{\Gamma_3} = 0$.

Therefore $\{B_1, B_2, B_3, B_4, b_1, b_2, b_3, a_3\}$ might be a subset of $G_{\Gamma_3,5}$ and we will extend it to a basis of $H_1(\Sigma_5; \mathbb{Z}_2)$ by adding two simple closed curves d_1, d_2 as in Figure 5 so that

$$\Gamma_3 = \Gamma(\{B_1, B_2, B_3, B_4, b_1, b_2, b_3, a_3, d_1, d_2\}).$$

Then Γ_3 is the graph as in Figure 7 and it satisfies Equations (4.6) and (4.7).




FIGURE 7. Graph Γ_3

Therefore $G_F(\Phi_{K_{0,1}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2) \leq G_{\Gamma_3,5}$ and, since $t_{c_2}, t_d \notin G_{\Gamma_3,5}$, we get

$$t_{c_2}, t_d \notin G_F(\Phi_{K_{0,1}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2)$$

and

$$t_{c_2}, t_d \notin G_F(\Phi_{K_{2p,2q+1}}(\eta_{1,2}^2) \cdot \eta_{1,2}^2)$$

for any $p, q \in \mathbb{Z}$.

Case 4: p and q are odd integers: We want to find a graph

$$\Gamma_4 = \Gamma(\{\gamma_1, \gamma_2, \dots, \gamma_{10}\})$$

satisfying

$$(4.9) \quad \chi_{\Gamma_4}(B_i) = \chi_{\Gamma_4}(\Phi_{K_{1,1}}(B_i)) = \chi_{\Gamma_4}(b_3) = \chi_{\Gamma_4}(b'_3) = \chi_{\Gamma_4}(a_3) = 1$$

for $i = 0, 1, 2, 3, 4, 5$ and

$$(4.10) \quad \chi_{\Gamma_4}(c_2) = \chi_{\Gamma_4}(d) = 0.$$


Note that we observe the following relations in $H_1(\Sigma_5; \mathbb{Z}_2)$:

$$(4.11) \quad \begin{array}{c|c|c} & \Phi_{K_{0,0}}(B_i) & \Phi_{K_{1,1}}(B_i) \\ \hline B_0 & B_0 + a_1 + b_1 + a_2 + b_2 & B_0 + a_1 + b_1 + a_2 + b_2 \\ \hline B_1 & B_1 + b_1 + b_2 + a_2 & B_1 + b_1 + a_2 + b_2 + c_2 + d \\ \hline B_2 & B_2 + a_1 + b_2 + a_2 & B_2 + a_1 + b_2 + a_2 + c_2 \\ \hline B_3 & B_3 + b_2 & B_3 + b_2 + c_2 \\ \hline B_4 & B_4 + a_2 & B_4 + a_2 + c_2 + d \\ \hline B_5 & B_5 & B_5 \end{array}$$

Hence, by Lemma 4.2 and Equations (4.9), (4.10) and (4.11), we need to find a graph Γ_4 such that

- even number of $\{a_1, b_1, a_2, b_2\}$ have $\chi_{\Gamma_4} = 0$
- even number of $\{b_1, b_2, a_2\}$ have $\chi_{\Gamma_4} = 0$
- odd number of $\{a_1, b_2, a_2\}$ have $\chi_{\Gamma_4} = 0$
- odd number of $\{b_2\}$ have $\chi_{\Gamma_4} = 0$

- $$\Gamma_4 = \Gamma(\{B_1, B_2, B_3, B_4, a_1, a_2, a_3, b_3, d_3, d_4\}).$$


$$\begin{aligned} \text{Order}(\psi_2(G_F(\xi_{p,q}))) &= 50030759116800, \\ \text{Order}(\langle \psi_2(G_F(\xi_{p,q}) \cup \{t_{c_2}\}) \rangle) &= 24815256521932800, \\ \text{Order}(\langle \psi_2(G_F(\xi_{p,q}) \cup \{t_d\}) \rangle) &= 24815256521932800, \\ \text{Order}(\psi_2(\mathcal{M}_5)) &= 24815256521932800, \end{aligned}$$

and it implies

$$t_{c_2}, t_d \notin G_F(\xi_{p,q}) \text{ for any } p, q \in \mathbb{Z}.$$

Theorem 4.5. $\xi_{p,q}$ is not equivalent to $\xi_{r,s}$ if $(p, q) \not\equiv (r, s) \pmod{2}$.

Proof. We obtain the following table from the proof of Lemma 4.3:

	$G_{\Gamma_i,5}$ does not contain
Γ_1	$t_{\Phi_{K_{1,0}}(B_j)}, (j = 1, 2, 3, 4), t_{\Phi_{K_{0,1}}(B_1)}, t_{\Phi_{K_{0,1}}(B_4)}, t_{\Phi_{K_{1,1}}(B_2)}, t_{\Phi_{K_{1,1}}(B_3)}$
Γ_2	$t_{\Phi_{K_{0,0}}(B_j)}, (j = 1, 2, 3, 4), t_{\Phi_{K_{0,1}}(B_2)}, t_{\Phi_{K_{0,1}}(B_3)}, t_{\Phi_{K_{1,1}}(B_1)}, t_{\Phi_{K_{1,1}}(B_4)}$
Γ_3	$t_{\Phi_{K_{0,0}}(B_1)}, t_{\Phi_{K_{0,0}}(B_4)}, t_{\Phi_{K_{1,0}}(B_2)}, t_{\Phi_{K_{1,0}}(B_3)}, t_{\Phi_{K_{1,1}}(B_j)}, (j = 1, 2, 3, 4)$
Γ_4	$t_{\Phi_{K_{0,0}}(B_2)}, t_{\Phi_{K_{0,0}}(B_3)}, t_{\Phi_{K_{1,0}}(B_1)}, t_{\Phi_{K_{1,0}}(B_4)}, t_{\Phi_{K_{0,1}}(B_j)}, (j = 1, 2, 3, 4)$

Therefore we get the following statement: $t_{\Phi_{K_{p,q}}(B_j)}$ is not contained in $G_{\Gamma_i,5}$ if and only if $t_{\Phi_{K_{\epsilon_p, \epsilon_q}}(B_j)}$ is not contained in $G_{\Gamma_i,5}$, where $\epsilon_p, \epsilon_q \in \{0, 1\}$ and $p \equiv \epsilon_p, q \equiv \epsilon_q \pmod{2}$. It implies that

$$\xi_{p,q} \not\sim \xi_{r,s} \text{ if } (p, q) \not\equiv (r, s) \pmod{2}.$$

For example, if $(p, q) \equiv (0, 0)$ and $(r, s) \equiv (1, 0) \pmod{2}$, then

$$t_{\Phi_{K_{p,q}}(B_j)} \notin G_{\Gamma_2,5}, \quad (j = 1, 2, 3, 4)$$

and $G_F(\xi_{r,s}) \leq G_{\Gamma_2,5}$. Therefore $t_{\Phi_{K_{p,q}}(B_j)} \in G_F(\xi_{p,q})$ but $t_{\Phi_{K_{p,q}}(B_j)} \notin G_F(\xi_{r,s})$ for $j = 1, 2, 3, 4$. It implies $G_F(\xi_{p,q}) \neq G_F(\xi_{r,s})$ and $\xi_{p,q} \not\sim \xi_{r,s}$. $\square$

Corollary 4.6. If $p \not\equiv q \pmod{2}$, then the knot surgery 4-manifold $E(2)_{K_{p,q}}$ has at least 2 nonisomorphic genus 5 Lefschetz fibration structures.

Proof. It follows from Lemma 3.1. Since $K_{p,q}$ is equivalent to $K_{q,p}$, we get a diffeomorphism $E(n)_{K_{p,q}} \approx E(n)_{K_{q,p}}$. Because $S^1 \times (S^3 \setminus \nu(K)) \approx S^1 \times (S^3 \setminus \nu(K^*))$, we also have $E(n)_K \approx E(n)_{K^*}$. So we get that $E(n)_{K_{p,q}} \approx E(n)_{K_{q,p}^*} \approx E(n)_{K_{-q,-p}}$. The last diffeomorphism is $E(n)_{K_{p,q}} \approx E(n)_{K_{q,p}} \approx E(n)_{K_{q,p}^*} \approx E(n)_{K_{-p,-q}}$. Therefore we have diffeomorphisms

$$E(2)_{K_{p,q}} \approx E(2)_{K_{q,p}} \approx E(2)_{K_{-p,-q}} \approx E(2)_{K_{-q,-p}}$$

and $\{\xi_{p,q}, \xi_{q,p}, \xi_{-p,-q}, \xi_{-q,-p}\}$ are Lefschetz fibration structures which have diffeomorphic underlying 4-manifolds. But, by Theorem 4.5, we know that $\xi_{p,q} \not\sim \xi_{q,p}$ because $(p, q) \not\equiv (q, p) \pmod{2}$. Therefore we get a conclusion. $\square$

Remark 4.7. At this time we do not know how to distinguish each elements in

$$\{ \xi_{p,q} \mid p, q \in \mathbb{Z} \}$$

up to Lefschetz fibration isomorphism.

Remark 4.8. We are interested in the question whether the knot surgery 4-manifold $E(2)_K$ admits infinitely many nonisomorphic Lefschetz fibrations over S^2 with the same generic fiber. In Theorem 3.4 we constructed a family of simply connected genus 5 Lefschetz fibrations over S^2 whose underlying space are all diffeomorphic and they are constructed from a pair of inequivalent prime fibred knots. We expect that they are strong candidates for admitting infinitely many nonisomorphic Lefschetz fibrations. We leave this problem for future research project.

REFERENCES

- [1] M. Akaho, *A connected sum of knots and Fintushel-Stern knot surgery on 4-manifolds*, Turkish J. Math. **30** (2006), no. 1, 87–93.
- [2] S. Akbulut, *Variations on Fintushel-Stern knot surgery on 4-manifolds*, Turkish J. Math. **26** (2002), no. 1, 81–92.
- [3] R. Fintushel and R. Stern, *Knots, links, and 4-manifolds*, Invent. Math. **134**(2), 363–400 (1998).
- [4] R. Fintushel and R. Stern, *Constructions of smooth 4-manifolds*, Proceedings of the International Congress of Mathematicians, Extra Vol. II (Berlin, 1998), 443–452.
- [5] R. Fintushel and R. Stern, *Families of simply connected 4-manifolds with the same Seiberg-Witten invariants*, Topology **43** (2004), no. 6, 1449–1467.
- [6] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4.12*, 2008.
- [7] R. Gompf and A. Stipsicz, *4-manifolds and Kirby calculus*, American Mathematical Society, Providence, RI, 1999.
- [8] Y. Gurtas, *Positive Dehn twist expressions for some new involutions in mapping class group*, 2004, arXiv:math.GT/0404310.
- [9] J. Harer, *How to construct all fibered knots and links*, Topology **21** (1982), no. 3, 263–280.
- [10] S. Humphries, e-mail communication.
- [11] S. Humphries, *Generators for the mapping class group*, Topology of low-dimensional manifolds, Lecture Notes in Math., vol. 722 (1979), Springer, 44–47.
- [12] T. Kanenobu, *Infinitely many knots with the same polynomial invariant*, Proc. Amer. Math. Soc. **97** (1986), no. 1, 158–162.
- [13] T. Kanenobu, *Examples on polynomial invariants of knots and links*, Math. Ann. **275** (1986), no. 4, 555–572.
- [14] A. Kas, *On the handlebody decomposition associated to a Lefschetz fibration*, Pacific J. Math. **89** (1980), no. 1, 89–104.
- [15] S. Kinoshita and H. Terasaka, *On unions of knots*, Osaka Math. J. **9** (1957), 131–153.
- [16] M. Korkmaz, *Noncomplex smooth 4-manifolds with Lefschetz fibrations*, Int. Math. Res. Notices **2001** (2001), no. 3, 115–128.
- [17] Y. Matsumoto, *Lefschetz fibrations of genus two—a topological approach*, Topology and Teichmüller spaces (Katinkulta, 1995), World Sci. Publishing, River Edge, NJ, 123–148.
- [18] J. Park and K.-H. Yun, *Nonisomorphic Lefschetz fibrations on knot surgery 4-manifolds*, Math. Ann. (2009), <http://dx.doi.org/10.1007/s00208-009-0366-0>.
- [19] *SAGE mathematics software, version 3.4*, <http://www.sagemath.org/>.
- [20] J. Stallings, *Constructions of fibred knots and links*, Algebraic and geometric topology, Part 2, Proc. Sympos. Pure Math., XXXII, Amer. Math. Soc., Providence, R.I., 55–60.
- [21] K.-H. Yun, *On the signature of a Lefschetz fibration coming from an involution*, Topology Appl. **153** (2006), no. 12, 1994–2012.
- [22] K.-H. Yun, *Twisted fiber sums of Fintushel-Stern’s knot surgery 4-manifolds*, Trans. Amer. Math. Soc. **360** (2008), no. 11, 5853–5868.

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, 599 GWANAK-RO, GWANAK-GU, SEOUL 151-747, REPUBLIC OF KOREA

E-mail address: jipark@snu.ac.kr

DEPARTMENT OF MATHEMATICS, SUNGSHIN WOMEN’S UNIVERSITY, 54-7 DONGSEONDONG-GIL, SEONGBUK-GU, SEOUL 136-742, REPUBLIC OF KOREA

E-mail address: kyun@sungshin.ac.kr